

DRAFT Medical Coverage Policy | Percutaneous Electrical Nerve Stimulation, Percutaneous Neuromodulation Therapy, and Restorative Neurostimulation Therapy



EFFECTIVE DATE: 02|01|2024

POLICY LAST REVIEWED: 10|04|2023

OVERVIEW

Percutaneous electrical nerve stimulation (PENS), percutaneous neuromodulation therapy (PNT), and restorative neurostimulation therapy (ReActiv8) combine the features of electroacupuncture and transcutaneous electrical nerve stimulation. Percutaneous electrical nerve stimulation is performed with needle electrodes while PNT uses very fine needle-like electrode arrays placed near the painful area to stimulate peripheral sensory nerves in the soft tissue. ReActiv8 is an implantable electrical neurostimulation system that stimulates the nerves that innervate the lumbar multifidus muscles.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

Medicare Advantage Plans

Restorative neurostimulation therapy (ReActiv8), Percutaneous electrical neurostimulation (PENS) or percutaneous neuromodulation therapy (PNT) for the treatment of chronic pain conditions is considered medically necessary if pain is effectively controlled by percutaneous stimulation and implantation of electrodes is warranted.

Note: Blue Cross & Blue Shield of Rhode Island (BCBSRI) must follow Centers for Medicare and Medicaid Services (CMS) guidelines, such as national coverage determinations or local coverage determinations for all Medicare Advantage Plans policies. Therefore, Medicare Advantage Plans policies may differ from Commercial products. In some instances, benefits for Medicare Advantage Plans may be greater than what is allowed by the CMS.

Commercial Products

Restorative neurostimulation therapy (ReActiv8), Percutaneous electrical neurostimulation (PENS) or percutaneous neuromodulation therapy (PNT) for the treatment of chronic pain conditions is considered not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes.

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable surgery or not medically necessary benefits/coverage.

BACKGROUND

A variety of chronic musculoskeletal or neuropathic pain conditions, including low back pain, neck pain, diabetic neuropathy, chronic headache, and surface hyperalgesia, present a substantial burden to patients, adversely affecting function and quality of life. Certain racial and ethnic groups are at a higher risk of developing diabetes, which may also put them at higher risk of developing complications from diabetes, such as diabetic neuropathy. According to a 2018 to 2019 National Health Interview Survey and data from the

Indian Health Service National Data Warehouse, American Indians and Alaska Natives had the highest reported rate of diagnosed diabetes at 14.5%. This was followed by 12.1% of Black individuals, 11.8% of Hispanic individuals, 9.5% of Asian individuals, and 7.4% of White individuals having diagnosed diabetes in 2018 or 2019.

These chronic pain conditions have typically failed other treatments, and percutaneous electrical nerve stimulation (PENS) and percutaneous neuromodulation therapy (PNT) have been evaluated as treatments to relieve unremitting pain.

Percutaneous electrical nerve stimulation is similar in concept to transcutaneous electrical nerve stimulation (TENS) (see evidence review 1.01.09) but differs in that needles are inserted either around or immediately adjacent to the nerves serving the painful area and are then stimulated. Percutaneous electrical nerve stimulation is generally reserved for patients who fail to get pain relief from TENS. Percutaneous electrical nerve stimulation is also distinguished from acupuncture with electrical stimulation. In electrical acupuncture, needles are also inserted just below the skin, but the placement of needles is based on specific theories regarding energy flow throughout the human body. In PENS, the location of stimulation is determined by proximity to the pain.

Percutaneous neuromodulation therapy is a variant of PENS in which fine filament electrode arrays are placed near the area causing pain. Some use the terms PENS and PNT interchangeably. It is proposed that PNT inhibits pain transmission by creating an electrical field that hyperpolarizes C fibers, thus preventing action potential propagation along the pain pathway.

The purpose of restorative neurostimulation therapy in individuals with chronic pain conditions is to provide a treatment option that is an alternative to or an improvement on existing therapies. The ReActiv8 System is an implantable electrical neurostimulation system that stimulates the nerves that innervate the lumbar multifidus muscles.

Medicare Advantage Plans

PENS, which involves stimulation of peripheral nerves by a needle electrode inserted through the skin is performed only in a physician's office, clinic, or hospital outpatient department. Therefore, it is covered only when performed by a physician or incident to physician's service. If pain is effectively controlled by percutaneous stimulation, implantation of electrodes is warranted.

Commercial Products

For individuals who have chronic pain conditions (e.g., back, neck, neuropathy, headache, hyperalgesia) who receive PENS, the evidence includes primarily small controlled trials and 2 systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and medication use. Two systematic reviews have not revealed consistent benefit from PENS in musculoskeletal pain disorders. One review concluded that PENS could decrease pain intensity but not related disability, while the other found no significant differences between PENS and TENS in mitigation of pain. These conclusions are uncertain due to important methodological limitations in individual trials included in these reviews, such as high heterogeneity with regard to application methods. In the highest quality trial of PENS conducted to date in chronic low back pain, no difference in outcomes was found between the active (30 minutes of stimulation with 10 needles) and the sham (5 minutes of stimulation with 2 needles) treatments. Smaller trials, which have reported positive results, are limited by unclear blinding and short-term follow-up. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have chronic pain conditions (eg, knee osteoarthritis) who receive PNT, the evidence consists of a randomized controlled trial (RCT). Relevant outcomes are symptoms, functional outcomes, quality of life, and medication use. The single trial is limited by lack of investigator blinding, unclear

participant blinding, and short-term follow-up. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have chronic pain conditions including low back pain who receive restorative neurostimulation therapy (ReActiv8), the evidence includes 1 sham-controlled RCT (N = 204), 1 prospective single-arm trial (N = 53), and a case series (N = 44). Relevant outcomes are symptoms, functional outcomes, quality of life, and medication use. In the RCT, there was no difference between groups on the primary endpoint of treatment response at 120 days, defined as the composite of 30% or greater reduction in VAS and no increase in pain medications (57.1% intervention vs 46.6% sham; p =.1377). Prespecified secondary analyses of primary outcome data favored the intervention group, but clinical significance is unclear. An uncontrolled follow-up phase of the RCT reported continued improvement in pain scores through 3 years but results are at high risk of bias due to lack of a control group and high attrition. Nonrandomized studies are limited by lack of blinding, no sham control, high attrition, and small sample sizes. Additional evidence from longer-term sham-controlled RCTs is needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

CODING

Medicare Advantage Plans and Commercial Products

There is not a specific code for PENS or PNT. Use the unlisted code below

64999 Unlisted procedure, nervous system

RELATED POLICIES

Lysis of Epidural Adhesions

Nerve Graft with Radical Prostatectomy

Occipital Nerve Stimulation – Insertion

Peripheral Subcutaneous Field Stimulation

Pulsed Radiofrequency for the Treatment of Chronic Pain

Sphenopalatine Ganglion Block for Headache

Unlisted Procedures

PUBLISHED

Provider Update, February 2024

Provider Update, December 2023

Provider Update, October 2022

Provider Update, October 2021

Provider Update, September 2020

REFERENCES

1. Centers for Disease Control and Prevention (CDC). By the Numbers: Diabetes in America. Updated March 2022; <https://www.cdc.gov/diabetes/health-equity/diabetes-by-the-numbers.html>. Accessed June 26, 2023.
2. Food & Drug Administration. 2020. ReActiv8 Implantable Neurostimulation System. Approval Order. https://www.accessdata.fda.gov/cdrh_docs/pdf19/P190021A.pdf. Accessed June 25, 2023.
3. Dworkin RH, Turk DC, Farrar JT, et al. Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain*. Jan 2005; 113(1-2): 9-19. PMID 15621359
4. Dworkin RH, Turk DC, Wyrwich KW, et al. Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. *J Pain*. Feb 2008; 9(2): 105-21. PMID 18055266
5. Gewandter JS, Dworkin RH, Turk DC, et al. Research design considerations for chronic pain prevention clinical trials: IMMPACT recommendations. *Pain*. Jul 2015; 156(7): 1184-1197. PMID 25887465
6. Plaza-Manzano G, Gómez-Chiguano GF, Cleland JA, et al. Effectiveness of percutaneous electrical nerve stimulation for musculoskeletal pain: A systematic review and meta-analysis. *Eur J Pain*. Jul 2020; 24(6): 1023-1044. PMID 32171035

7. Beltran-Alacreu H, Serrano-Muñoz D, Martín-Caro Álvarez D, et al. Percutaneous Versus Transcutaneous Electrical Nerve Stimulation for the Treatment of Musculoskeletal Pain. A Systematic Review and Meta-Analysis. *Pain Med.* Aug 01 2022; 23(8): 1387-1400. PMID 35167691
8. Ghoname EA, Craig WF, White PF, et al. Percutaneous electrical nerve stimulation for low back pain: a randomized crossover study. *JAMA.* Mar 03 1999; 281(9): 818-23. PMID 10071003
9. Ghoname ES, Craig WF, White PF, et al. The effect of stimulus frequency on the analgesic response to percutaneous electrical nerve stimulation in patients with chronic low back pain. *Anesth Analg.* Apr 1999; 88(4): 841-6. PMID 10195535
10. Hamza MA, Ghoname EA, White PF, et al. Effect of the duration of electrical stimulation on the analgesic response in patients with low back pain. *Anesthesiology.* Dec 1999; 91(6): 1622-7. PMID 1059860224.
- Medeiros LF, Caumo W, Dussán-Sarria J, et al. Effect of Deep Intramuscular Stimulation and Transcranial Magnetic Stimulation on Neurophysiological Biomarkers in Chronic Myofascial Pain Syndrome. *Pain Med.* Jan 2016; 17(1): 122-35. PMID 26408420
11. Weiner DK, Rudy TE, Glick RM, et al. Efficacy of percutaneous electrical nerve stimulation for the treatment of chronic low back pain in older adults. *J Am Geriatr Soc.* May 2003; 51(5): 599-608. PMID 12752833
12. Topuz O, Ozfidan E, Ozgen M, Ardic F. Efficacy of transcutaneous electrical nerve stimulation and percutaneous neuromodulation therapy in chronic low back pain. *J Back Musculoskeletal Rehabil.* 2004; 17:127-133.
13. Yokoyama M, Sun X, Oku S, et al. Comparison of percutaneous electrical nerve stimulation with transcutaneous electrical nerve stimulation for long-term pain relief in patients with chronic low back pain. *Anesth Analg.* Jun 2004; 98(6): 1552-1556. PMID 15155304
14. Weiner DK, Perera S, Rudy TE, et al. Efficacy of percutaneous electrical nerve stimulation and therapeutic exercise for older adults with chronic low back pain: a randomized controlled trial. *Pain.* Nov 30 2008; 140(2): 344-357. PMID 18930352
15. Perez-Palomares S, Oliván-Blázquez B, Magallon-Botaya, et al. Percutaneous electrical nerve stimulation versus dry needling: effectiveness in the treatment of chronic low back pain. *J Musculoskeletal Pain.* 2010; 18:23-30.
16. Weiner DK, Rudy TE, Morone N, et al. Efficacy of periosteal stimulation therapy for the treatment of osteoarthritis-associated chronic knee pain: an initial controlled clinical trial. *J Am Geriatr Soc.* Oct 2007; 55(10): 1541-7. PMID 17908057
17. Weiner DK, Moore CG, Morone NE, et al. Efficacy of periosteal stimulation for chronic pain associated with advanced knee osteoarthritis: a randomized, controlled clinical trial. *Clin Ther.* Nov 2013; 35(11):1703-20.e5. PMID 24184053
18. da Graca-Tarragó M, Deitos A, Patrícia Brietzke A, et al. Electrical Intramuscular Stimulation in Osteoarthritis Enhances the Inhibitory Systems in Pain Processing at Cortical and Cortical Spinal System. *Pain Med.* May 01 2016; 17(5): 877-891. PMID 26398594
19. Elbadawy MA. Effectiveness of Periosteal Stimulation Therapy and Home Exercise Program in the Rehabilitation of Patients With Advanced Knee Osteoarthritis. *Clin J Pain.* Mar 2017; 33(3): 254-263. PMID 27513639
20. Dunning J, Butts R, Henry N, et al. Electrical dry needling as an adjunct to exercise, manual therapy and ultrasound for plantar fasciitis: A multi-center randomized clinical trial. *PLoS One.* 2018; 13(10):e0205405. PMID 30379937
21. da Graca-Tarragó M, Lech M, Angoleri LDM, et al. Intramuscular electrical stimulus potentiates motor cortex modulation effects on pain and descending inhibitory systems in knee osteoarthritis: a randomized, factorial, sham-controlled study. *J Pain Res.* 2019; 12: 209-221. PMID 30655690
22. León-Hernández JV, Martín-Pintado-Zugasti A, Frutos LG, et al. Immediate and short-term effects of the combination of dry needling and percutaneous TENS on post-needling soreness in patients with chronic myofascial neck pain. *Braz J Phys Ther.* Jul 11 2016; 20(5): 422-431. PMID 27410163
23. Sumen A, Sarsan A, Alkan H, et al. Efficacy of low level laser therapy and intramuscular electrical stimulation on myofascial pain syndrome. *J Back Musculoskeletal Rehabil.* 2015; 28(1): 153-8. PMID 25061034

24. Medeiros LF, Caumo W, Dussán-Sarria J, et al. Effect of Deep Intramuscular Stimulation and Transcranial Magnetic Stimulation on Neurophysiological Biomarkers in Chronic Myofascial Pain Syndrome. *Pain Med.* Jan 2016; 17(1): 122-35. PMID 26408420
25. Botelho L, Angoleri L, Zortea M, et al. Insights About the Neuroplasticity State on the Effect of Intramuscular Electrical Stimulation in Pain and Disability Associated With Chronic Myofascial Pain Syndrome (MPS): A Double-Blind, Randomized, Sham-Controlled Trial. *Front Hum Neurosci.* 2018; 12:388. PMID 30459575
26. Dunning J, Butts R, Young I, et al. Periosteal Electrical Dry Needling as an Adjunct to Exercise and Manual Therapy for Knee Osteoarthritis: A Multicenter Randomized Clinical Trial. *Clin J Pain.* Dec 2018;34(12): 1149-1158. PMID 29864043
27. Yoshimizu M, Teo AR, Ando M, Kiyohara K, Kawamura T. Relief of chronic shoulder and neck pain by electro-acupuncture and transcutaneous electrical nervous stimulation: A randomized crossover trial. *Med Acupunct* 2012;24(2):97103.
28. Ng MM, Leung MC, Poon DM. The effects of electro-acupuncture and transcutaneous electrical nerve stimulation on patients with painful osteoarthritic knees: a randomized controlled trial with follow-up evaluation. *J Altern Complement Med.* Oct 2003; 9(5): 641-9. PMID 14629842
29. Tsukayama H, Yamashita H, Amagai H, et al. Randomised controlled trial comparing the effectiveness of electroacupuncture and TENS for low back pain: a preliminary study for a pragmatic trial. *Acupunct Med.* Dec 2002; 20(4): 175-80. PMID 12512791
30. Cheng RSS, Pomeranz B. Electrotherapy of chronic musculoskeletal pain: Comparison of electroacupuncture and acupuncture-like transcutaneous electrical nerve stimulation. *Cochrane Library. Clin J Pain* 1986;2(3):1439.
31. Lehmann TR, Russell DW, Spratt KF, et al. Efficacy of electroacupuncture and TENS in the rehabilitation of chronic low back pain patients. *Pain.* Sep 1986; 26(3): 277-290. PMID 2946016
32. Ghoname EA, White PF, Ahmed HE, et al. Percutaneous electrical nerve stimulation: an alternative to TENS in the management of sciatica. *Pain.* Nov 1999; 83(2): 193-9. PMID 10534590
33. White PF, Craig WF, Vakharia AS, et al. Percutaneous neuromodulation therapy: does the location of electrical stimulation effect the acute analgesic response?. *Anesth Analg.* Oct 2000; 91(4): 949-54. PMID 11004055
34. Hamza MA, White PF, Craig WF, et al. Percutaneous electrical nerve stimulation: a novel analgesic therapy for diabetic neuropathic pain. *Diabetes Care.* Mar 2000; 23(3): 365-70. PMID 10868867
35. Ahmed HE, White PF, Craig WF, et al. Use of percutaneous electrical nerve stimulation (PENS) in the short-term management of headache. *Headache.* Apr 2000; 40(4): 311-5. PMID 10759936
36. Raphael JH, Raheem TA, Southall JL, et al. Randomized double-blind sham-controlled crossover study of short-term effect of percutaneous electrical nerve stimulation in neuropathic pain. *Pain Med.* Oct 2011; 12(10): 1515-22. PMID 21883874
37. Kang RW, Lewis PB, Kramer A, et al. Prospective randomized single-blinded controlled clinical trial of percutaneous neuromodulation pain therapy device versus sham for the osteoarthritic knee: a pilot study. *Orthopedics.* Jun 2007; 30(6): 439-45. PMID 17598487
38. Gilligan C, Volschenk W, Russo M, et al. An implantable restorative-neurostimulator for refractory mechanical chronic low back pain: a randomized sham-controlled clinical trial. *Pain.* Oct 2021;162(10): 2486-2498. PMID 34534176
39. Food & Drug Administration. 2020. ReActiv8 Implantable Neurostimulation System: Summary of Safety and Effectiveness Data. https://www.accessdata.fda.gov/cdrh_docs/pdf19/P190021B.pdf. Accessed June 23, 2023.
40. Gilligan C, Volschenk W, Russo M, et al. Long-Term Outcomes of Restorative Neurostimulation in Patients With Refractory Chronic Low Back Pain Secondary to Multifidus Dysfunction: Two-Year Results of the ReActiv8-B Pivotal Trial. *Neuromodulation.* Jan 2023; 26(1): 87-97. PMID 35088722
41. Gilligan C, Volschenk W, Russo M, et al. Three-Year Durability of Restorative Neurostimulation Effectiveness in Patients With Chronic Low Back Pain and Multifidus Muscle Dysfunction. *Neuromodulation.* Jan 2023; 26(1): 98-108. PMID 36175320

